

# Random Walks and Variance Ratio Tests

**Christopher Ting**

<http://cting.x10host.com/TMU/TMU.html>

<http://www.mysmu.edu/faculty/christophert/>

✉: [christophert@smu.edu.sg](mailto:christophert@smu.edu.sg)

June 7, 2018

# Broad Lesson Plan

- 1 Introduction
- 2 Law of Iterated Expectations
- 3 Random Walks
- 4 Chi-Square R.V
- 5 Variance Ratio Test
- 6 Takeaways

## Four Big Questions

### (1) Are stock returns predictable?

Common sense answer: No. Otherwise, everyone will make money without risk.

### (2) Are asset returns 100% random?

Investment firms: No. Otherwise, we will have no client!

### (3) Will stock market indexes move up over 30 years?

Common sense answer: Yes. Otherwise why would anyone invest in risky stocks?

### (4) Can mutual funds beat their benchmark stock market indexes?

Mutual fund managers: Yes. It's our job to beat the market.

Academic professors: No. On average, mutual funds are equally likely to beat the market; after costs, the alpha is negative.

# Types of Market Efficiency

- ✿ Weak form efficiency: information in past returns or prices
- ✿ Semi-strong form efficiency: all public information
- ✿ Strong form efficiency: all private information (fundamental research on companies, economies, etc)

## A Paradox

If the market is (strong form) efficient, no one will bother to gather private information because they can't beat the market. But if no one gathers private information, then the market cannot possibly reflect any private information!

John Y. Campbell

### On the Impossibility of Informationally Efficient Markets

Strong form efficient market hypothesis is theoretically implausible, unless information is free. If information is costly to gather (i.e., costly to do fundamental research on companies), then investors will only gather information if the benefit outweighs the cost.

Grossman and Stiglitz

## Implications in Practice

- ☀ Reality: Research is still ongoing to obtain and analyze new private information.
- ☀ Consequence: Research results will push prices to some extent, i.e., prices will reflect some but not all of the new information.
- ☀ In equilibrium, the marginal benefit of obtaining new information will be equal to the marginal cost of doing research.
- ☀ What are the practical implications?
  - ☀ You can beat the market, as long as your cost of information acquisition and analysis is less than other people's.
  - ☀ If you have no particular advantage, then you should hold primarily low-cost index funds (and free ride on the information obtained by others), possibly with a tilt towards value.

# Martingale

## Definition 1

Suppose  $\{P_t\}_{t=1}^T$  is a **stochastic process**. It is said to be a **martingale** if

$$\mathbb{E}(P_t | P_{t-1}, P_{t-2}, \dots) = P_{t-1}.$$

Equivalently, since  $P_{t-1}$  is a known constant at time  $t - 1$ ,

$$\mathbb{E}(P_t - P_{t-1} | P_{t-1}, P_{t-2}, \dots) = 0.$$

## Implications

- A** Given the information  $\{P_{t-i}\}_{i=1}^t$  up to time  $t - 1$ , the **best forecast** of  $P_t$  is  $P_{t-1}$ .
- B** Obviously, in most cases,  $P_t - P_{t-1} \neq 0$  and in that sense, the forecast is wrong and thus  $P_t$  and return are **unpredictable**.

# Fundamental Value

## Definition 2

The conditional expectation of a random variable  $X$  is a probability-weighted mean of the different possible values of  $X$

$$\mathbb{E}(X|\mathcal{I}) = \sum_{i=1}^n \pi_i X_i,$$

where the probability  $\pi_i$  for each  $i$  is conditional on the information set  $\mathcal{I}$ .

## Remark

The expected value conditional on the information set  $\mathcal{I}$  is the **rational expectation** of an asset's payoff, i.e.,  $\mathbb{E}(X|\mathcal{I})$  is the **fundamental value** of the asset.

## A Very Important Question

If returns are unpredictable, does it mean that the market price is equal to the fundamental value?

# Law of Iterated Expectations (LIE)

## Proposition 1

For two random variables  $X$  and  $Y$ , the law of iterated expectations says that

$$\mathbb{E}_Y \left( \mathbb{E}_{X|Y} (X | Y) \right) = \mathbb{E} (X).$$

## Example

Suppose a portfolio has two stocks,  $A$  and  $B$ . Stock  $A$  ( $B$ ) has a mean return of 6% (15%) per annum. The investment in Stock  $A$  ( $B$ ) is 2 (4) billion dollars. What is the portfolio's expected return?

Answer: Based on the investment amounts, the weight of  $A$  is  $1/3$  and that of  $B$  is  $2/3$ . The return  $r_P$  of the portfolio is

$$\begin{aligned} \mathbb{E}(r_P) &= \mathbb{E}(r_P|A) \mathbb{P}(A) + \mathbb{E}(r_P|B) \mathbb{P}(B) \\ &= 6\% \times \frac{1}{3} + 15\% \times \frac{2}{3} = 12\% \end{aligned}$$

# Proof of LIE

$$\begin{aligned}
 \mathbb{E}_Y \left( \mathbb{E}_{X|Y} (X | Y) \right) &= \mathbb{E}_Y \left[ \sum_x x \cdot \mathbb{P}(X = x | Y) \right] \\
 &= \sum_y \left[ \sum_x x \cdot \mathbb{P}(X = x | Y = y) \right] \cdot \mathbb{P}(Y = y) \\
 &= \sum_y \sum_x x \cdot \mathbb{P}(X = x | Y = y) \cdot \mathbb{P}(Y = y) \\
 &= \sum_x x \sum_y \mathbb{P}(X = x | Y = y) \cdot \mathbb{P}(Y = y) \\
 &= \sum_x x \sum_y \mathbb{P}(X = x, Y = y) \\
 &= \sum_x x \cdot \mathbb{P}(X = x)
 \end{aligned}$$

# General LIE

## Proposition 2

For a **probability space**  $(\Omega, \mathcal{F}_\infty, \mathbb{P})$ , suppose the  **$\sigma$  algebras** are such that  $\mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \mathcal{F}_\infty$ . A random variable  $X$  is defined on this space. The law of iterated expectations states that

$$\mathbb{E} \left( \mathbb{E} (X | \mathcal{F}_2) \middle| \mathcal{F}_1 \right) = \mathbb{E} (X | \mathcal{F}_1).$$

✱ Note that

$$\begin{aligned} \int_{F_1} \mathbb{E} \left( \mathbb{E} (X | \mathcal{F}_2) \middle| \mathcal{F}_1 \right) d\mathbb{P} &= \int_{F_1} \mathbb{E} (X | \mathcal{F}_2) d\mathbb{P} \\ &= \int_{F_1} X d\mathbb{P} \text{ holds for all } F_1 \in \mathcal{F}_1 \subseteq \mathcal{F}_2 \end{aligned}$$

✱ In the special case of  $\mathcal{F}_1 = \{\emptyset, \Omega\}$  and  $\mathcal{F}_2 = \sigma(Y)$ , the  $\sigma$  algebra generated by random variable  $Y$ , we have

$$\mathbb{E}(\mathbb{E}(X|Y)) = \mathbb{E}(X).$$

## LIE over Time

\* Suppose time  $t$  information is less than time  $t + 1$  information, which is less than time  $t + 2$  information.

\* Let

$$\mathbb{E}_t(X) := \mathbb{E}(X | \text{Time } t \text{ information})$$

\* By the Law of Iterated Expectations,


$$\mathbb{E}_t(X) = \mathbb{E}_t(\mathbb{E}_{t+1}(X))$$

\* Alternatively,

$$\mathbb{E}_t(\mathbb{E}_{t+1}(X) - X) = 0. \quad (1)$$

\* The expectation at  $t$  of the **mistake**  $\mathbb{E}_{t+1}(X) - X$  you are going to make at  $t + 1$  is zero.

# Example: Tossing Two Fair Coins



## Example: Tossing Two Fair Coins (Cont'd)

- \* Let  $X$  be the number of heads. Then, the direct method obtains

$$\mathbb{E}_t(X) = \frac{1}{4} \times 2 + \frac{1}{2} \times 1 + \frac{1}{4} \times 0 = 1$$

- \* The LIE's method obtains

$$\begin{aligned} \mathbb{E}_{t+1}(X) &= \frac{3}{2} \quad \text{or} \quad \frac{1}{2} \\ \mathbb{E}_t(\mathbb{E}_{t+1}(X)) &= \frac{1}{2} \times \frac{3}{2} + \frac{1}{2} \times \frac{1}{2} = 1. \end{aligned}$$

- \* Hence, we have illustrated that

$$\mathbb{E}_t(X) = \mathbb{E}_t(\mathbb{E}_{t+1}(X)).$$

## Verification of Alternative Formalism (1)

- \* With respect to  $t$ , the future forecast is  $\mathbb{E}_{t+1}(X)$  for  $X$ 's value at  $t + 2$ .
- \* The forecast made at  $t + 1$  is either  $1/2$  or  $3/2$ .
- \* In the case of  $1/2$ , the correct value is either  $0$  or  $1$  at  $t + 2$  with equal probability. So the error is either  $(1/2 - 0)$  or  $(1/2 - 1)$ .
- \* In the case of  $3/2$ , the correct value is either  $1$  or  $2$  at  $t + 2$  with equal probability. So the error is either  $(3/2 - 1)$  or  $(3/2 - 2)$ .
- \* Based on the probability measure at  $t$ ,

$$\begin{aligned} \mathbb{E}_t(\mathbb{E}_{t+1}(X) - X) &= \left(\frac{1}{4} \times \left(\frac{1}{2} - 0\right)\right) + \left(\frac{1}{4} \times \left(\frac{1}{2} - 1\right)\right) \\ &\quad + \left(\frac{1}{4} \times \left(\frac{3}{2} - 1\right)\right) + \left(\frac{1}{4} \times \left(\frac{3}{2} - 2\right)\right) = 0. \end{aligned}$$

# Fundamental Value and Unpredictable Returns

- \* Suppose prices are based on the fundamental value of the future payoff  $X$ , i.e.,  $P_t = \mathbb{E}_t(X)$  and  $P_{t+1} = \mathbb{E}_{t+1}(X)$
- \* By the LIE, since  $\mathbb{E}_t(X) = \mathbb{E}_t(\mathbb{E}_{t+1}(X))$ , the price change is

$$\begin{aligned}
 \Delta P_{t+1} &:= P_{t+1} - P_t \\
 &= P_{t+1} - \mathbb{E}_t(X) \\
 &= P_{t+1} - \mathbb{E}_t(\mathbb{E}_{t+1}(X)) \\
 &= P_{t+1} - \mathbb{E}_t(P_{t+1}).
 \end{aligned}$$

- \* Therefore, price change (and hence return) is equal to the “mistake” or error  $P_{t+1} - \mathbb{E}_t(P_{t+1})$ , which is unpredictable.

# Fundamental Value and Rational Prediction

\* Now, by definition,

$$\Delta P_{t+1} = \mathbb{E}_{t+1}(X) - \mathbb{E}_t(X).$$

= future rational expectation – current rational expectation

\* Implications

- ✪ Price change or return is equally likely to be positive and negative, just like tossing a fair coin.
- ✪ Therefore, if you are rational, you cannot predict how you will change your mind in the future!

\* Insight

LIE connects fundamental value with unpredictable returns.

# Random Walk 1

## Definition 3

Random walk  $\{P_t\}_{t=1}^T$ , as a stochastic process, is expressed as

$$P_t = \mu + P_{t-1} + \epsilon_t, \quad \epsilon_t \sim \text{IID}(0, \sigma^2),$$

where

$\mu$  rate of drift

$\epsilon_t$  random "noise"

$\sigma^2$  variance

$\text{IID}(0, \sigma^2)$  **i**ndependently and **i**dentically **d**istributed  
with mean 0 and variance  $\sigma^2$ .


★ Note that no explicit distribution is specified for  $\epsilon_t$ .

# Properties of Random Walk 1

\* Let  $P_0$  be the initial position of the random walk at time 0. At time  $t$ ,

$$P_t = P_0 + \mu t + \sum_{s=0}^{t-1} \epsilon_{t-s}. \quad (2)$$

\* Suppose  $\epsilon_t$  is either 1 or -1, with  $\mathbb{P}(\epsilon_t = 1) = 0.5 = \mathbb{P}(\epsilon_t = -1)$ .



```
import numpy as np
import matplotlib.pyplot as plt

up_probability, n, paths = 0.5, 100, 20

np.random.seed(12)
p, r = np.zeros((n+1,paths),dtype=float), np.zeros((n+1,paths),dtype=float)
for z in range(paths):
    r[:, z] = np.random.random(n+1)

t = range(n+1)
for z in range(paths):
    for i in t[1:n+1]:
        if r[i, z] > up_probability:
            p[i,z] = p[i-1,z]+1
        else:
            p[i,z] = p[i-1,z]-1
    plt.plot(t, p[:,z])

plt.grid()
title = str(paths) + ' Simulated Random Walks'
plt.title(title)
plt.autoscale(enable=True, axis='x', tight=True)
plt.savefig('rw.png', format = 'png', dpi=288)
plt.show()
```

# Concept Checker

\* For the random walk  $P_t = \mu + P_{t-1} + \epsilon_t$  to be a martingale, you need

A.  $\mu = 0$

B.  $\mathbb{E}(\epsilon_t) = 0$

C.  $P_0 = 0$

D. None of the above, because random walk cannot be a martingale

# Linear Scaling Law

- \* Rewriting (2), we obtain


$$P_t - P_0 = \mu t + \sum_{s=0}^{t-1} \epsilon_{t-s}.$$

- \* It is easy to find that  $\mathbb{V}(\epsilon_t) = 1$  for the assumption in Slide 20.
- \* Therefore, since  $\mathbb{C}(\epsilon_t, \epsilon_s) = 0$  for  $t \neq s$ ,

$$\mathbb{V}(P_t - P_0) = \mathbb{V}\left(\sum_{s=0}^{t-1} \epsilon_{t-s}\right) = \sum_{s=0}^{t-1} \mathbb{V}(\epsilon_{t-s}) = t$$

- \* The variance from the origin  $P_0$  scales linearly with time  $t$  as it increases.

# Simulation of Linear Scaling Law



# More General Random Walks

## Random Walk 2

The random noise  $\epsilon_t$  is still independent but not identically distributed. In particular, the variance  $\sigma_t^2$  of the noise is different at different  $t$ .

## Random Walk 3

The random noise  $\epsilon_t$  is still identically distributed but the independence assumption is relaxed to zero covariance, i.e.,

$$\mathbb{C}(\epsilon_t, \epsilon_{t-s}) = 0.$$

for  $s \neq 0$ . In other words, the random noise does not have any linear correlation structure. However, nonlinear correlation, for example,  $\mathbb{C}(\epsilon_t^2, \epsilon_{t-s}^2) \neq 0$  is not assumed,

## Random Walk 4

The most general: neither independent nor identically distributed

## A Problem with Arithmetic Random Walk

\* The price of an asset such as stock is positive. But the arithmetic random walk 1 defined in Slide 19 can lead to a negative price if  $\epsilon_t$  is very negative.

\* Therefore, consider a random variable  $\xi_{t+1} > 0$  such that  $\mathbb{E}_t(\xi_{t+1}) = 1$  and

$$P_t = P_{t-1}\xi_t.$$

\* Then

$$\mathbb{E}_{t-1}(P_t) = P_{t-1}.$$

\* **In this model, log return  $r_t$  is noise!**

$$\begin{aligned} r_t &= \ln P_t - \ln P_{t-1} \\ &= \ln P_{t-1} + \ln \xi_t - \ln P_{t-1} \\ &= \ln \xi_t =: u_t \end{aligned}$$

# Geometric Random Walk

## Definition 4

Geometric random walk with drift  $\mu$  is defined as

$$\ln P_t = \mu + \ln P_{t-1} + u_t,$$

where  $u_t$  is not correlated over time and identically distributed with mean 0 and  $\mathbb{V}(u_t) = \sigma^2$ .

➤ In other word, two assumptions are made or the log return  $r_t$

✳ **Zero covariance:**  $\mathbb{C}(r_s, r_t) = 0$  for any  $s \neq t$


✳ **Homoskedasticity:**  $\mathbb{V}(r_t) = \sigma^2$

➤ Definition 4 implies that

$$r_t := \ln P_t - \ln P_{t-1} = \mu + u_t. \quad (3)$$

# Real-World Example

CK Hutchison Holdings Limited



## Variance of a Sum of $q$ Daily Log Returns


- When the daily log return  $r_t$  is treated as a random variable, the variance of a sum of  $q$  daily log returns is

$$\mathbb{V}\left(\sum_{t=1}^q r_t\right) = \sum_{t=1}^q \mathbb{V}(r_t) + 2 \sum_{t=1}^q \sum_{s < t} \mathbb{C}(r_s, r_t).$$

- Under the two assumptions in Slide 27,

$$\mathbb{V}(r_{t(q)}) := \mathbb{V}\left(\sum_{t=1}^q r_t\right) = q\sigma^2.$$

# Random Walk → Diffusion



## Starting Point of Chi-Square Random Variable

✦ Suppose  $Z$  is a standard normal random variable, i.e.,  $Z \sim N(0, 1)$ . Then  $Z^2$  is a chi-square random variable with one degree of freedom.

✦ The expected value of  $Z^2$  is, since  $Z \sim N(0, 1)$ ,

$$\mathbb{E}(Z^2) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} z^2 e^{-\frac{z^2}{2}} dz = 1.$$

✦ By the definition of variance,

$$\mathbb{V}(Z^2) = \mathbb{E}(Z^4) - (\mathbb{E}(Z^2))^2.$$

✦ We need to show that

$$\mathbb{E}(Z^4) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} z^4 e^{-\frac{z^2}{2}} dz = 3.$$

# Kurtosis

$$\begin{aligned}
 \int_{-\infty}^{\infty} z^4 e^{-z^2/2} dz &= - \int_{-\infty}^{\infty} z^3 de^{-z^2/2} \\
 &= - z^3 e^{-z^2/2} \Big|_{-\infty}^{+\infty} + 3 \int_{-\infty}^{\infty} z^2 e^{-z^2/2} dz \\
 &= -3 \int_{-\infty}^{\infty} z de^{-z^2/2} \\
 &= -3z e^{-z^2/2} \Big|_{-\infty}^{+\infty} + 3 \int_{-\infty}^{\infty} e^{-z^2/2} dz = 3\sqrt{2\pi}
 \end{aligned}$$

Therefore,

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} z^4 e^{-z^2/2} dz = 3$$

➤ It follows that the variance of  $Z^2$  is

$$\mathbb{V}(Z^2) = 3 - 1^2 = 2.$$

## Chi-Square Random Variable with $q$ Degrees

⇒ In general, if  $Z_1, \dots, Z_q$  are independent, standard normal random variables, then the sum of their squares,

$$Y = \sum_{i=1}^q Z_i^2,$$

is distributed according to the chi-squared distribution with  $q$  degrees of freedom, i.e.,  $Y \sim \chi_q^2$ .

⇒ By the assumption of independence,

$$\mathbb{V}(Y) = \sum_{i=1}^q \mathbb{V}(Z_i^2) = \sum_{i=1}^q 2 = 2q.$$

# Variance Ratio

## Definition 5

The **variance ratio** is defined as

$$\text{VR}(q) := \frac{\mathbb{V}(r_t(q))}{q\sigma^2},$$

- $\text{VR}(q)$  should be equal to 1 when the conditions of log returns being **serially uncorrelated** and **homoskedastic** are satisfied.
- The variance ratio test is a test of

$$H_0 : \text{VR}(q) - 1 = 0 \quad \text{versus} \quad H_1 : \text{VR}(q) - 1 \neq 0$$

- If the null hypothesis cannot be rejected, then it means that the two assumptions are consistent with the reality. Conversely, a rejection of  $H_0$  implies that at least one of the two assumptions is inconsistent with reality.

## Sample Mean and Variance of Daily Log Returns

- To set up the framework for inference, we recall a few definitions and facts. The sample mean of daily log returns is estimated as usual,

$$\hat{r}_1 = \frac{1}{T} \sum_{t=1}^T r_t. \quad (4)$$

- But the sample variance of daily log returns is instead estimated as

$$\hat{\sigma}_1^2 = \frac{1}{T} \sum_{t=1}^T (r_t - \hat{r}_1)^2. \quad (5)$$

- The subscript of 1 in  $\hat{r}_1$  and  $\hat{\sigma}_1^2$  is meant to indicate that these estimates are for daily log returns.

## Distribution of Daily Variance Estimate

□ By the law of large numbers, as  $T \rightarrow \infty$ ,

$$\mathbb{E}(\hat{\sigma}_1^2) = \frac{1}{T} \sum_{t=1}^T \mathbb{E}\left((r_t - \hat{r}_1)^2\right) \rightarrow \sigma^2$$

□ When  $T$  is very large, asymptotically, and by the central limit theorem, and since  $r_t = \mu + u_t$  as in Equation (3)

$$\begin{aligned} \mathbb{V}(\hat{\sigma}_1^2) &= \frac{1}{T^2} \sum_{t=1}^T \mathbb{V}\left((r_t - \hat{r}_1)^2\right) = \frac{1}{T^2} \sum_{t=1}^T \mathbb{V}\left((\mu + u_t - \hat{r}_1)^2\right) \\ &\rightarrow \frac{1}{T} \mathbb{V}(u_t^2) = \frac{1}{T} \mathbb{V}(\sigma^2 Z^2) = \frac{\sigma^4}{T} \mathbb{V}(Z^2) = \frac{2\sigma^4}{T}, \end{aligned}$$

where  $Z \sim N(0, 1)$ , and  $\mathbb{V}(Z^2)$  is the variance of the chi-square random variable with 1 degree of freedom ( $\chi_1^2$ ), which equals 2.

## Summary for Daily Sample Variance Estimate

- By the central limit theorem, as  $T$  becomes larger and larger,

$$\hat{\sigma}_1^2 \sim N\left(\sigma^2, \frac{2\sigma^4}{T}\right)$$

- Equivalently,

$$\sqrt{T}(\hat{\sigma}_1^2 - \sigma^2) \sim N(0, 2\sigma^4),$$

or

$$\sqrt{T}\left(\frac{\hat{\sigma}_1^2}{\sigma^2} - 1\right) \sim N(0, 2),$$

## Multi-Period Mean and Variance

- An important property of log return is the **telescoping sum**:

$$\begin{aligned} \ln \left( \frac{P_t}{P_0} \right) &= \ln(P_t) - \ln(P_{t-1}) + \ln P_{t-1} - \ln P_{t-2} + \ln P_{t-2} \\ &\quad + \cdots + \ln P_2 - \ln P_1 + \ln P_1 - \ln P_0 \end{aligned}$$

- Daily log return  $r_s := \ln P_s - \ln P_{s-1}$  for  $s = 1, 2, \dots, t$ ,

$$\ln P_t - \ln P_0 = \sum_{s=1}^t r_s$$

- Hence for log return

$$\begin{aligned} \mathbb{E} \left( \ln \left( \frac{P_t}{P_0} \right) \right) &= \sum_{s=1}^t \mathbb{E}(r_s) = t\mu \\ \mathbb{V} \left( \ln \left( \frac{P_t}{P_0} \right) \right) &= \sum_{s=1}^t \mathbb{V}(r_s) = t\sigma^2 \end{aligned}$$

## Concept Checker

♣ Sharpe ratio is a measure of risk-adjusted return:

$$R_{\text{Sharpe}} = \frac{\mathbb{E}(r_{i,t} - r_{f,t})}{\sqrt{\mathbb{V}(r_{i,t})}}$$

♣ You use the daily frequency to estimate the average excess log return of portfolio  $i$  and the standard deviation of the log return on portfolio  $i$ . How should you annualize the Sharpe ratio?

♣ Specifically, suppose the average excess return is 0.02% and the daily volatility is 1.1%. Suppose each year has 252 trading days. What is the annualized Sharpe ratio?

## Estimation of $q$ -Daily Sample Mean and Variance

- The  $q$ -daily return is

$$r(qj) := \ln P_{qj} - \ln P_{q(j-1)} = r_{qj} + r_{qj-1} + \cdots + r_{qj-(q-1)} + r_{q(j-1)}, \quad (6)$$

- for  $j = 1, 2, \dots, M$ , where  $M$  is the maximum number of **non-overlapping**  $q$ -daily returns that are obtainable from  $T + 1$  prices starting from  $P_0$ .

- As a matter of fact,  $M = \left\lfloor \frac{T}{q} \right\rfloor$ .

- The sample average of  $r(qj)$  is  $q$  times of  $\hat{r}_1$  (sample estimate of  $\mu$ ), i.e.,  $q\hat{r}_1$ .

- The sample variance is estimated as

$$\hat{\sigma}_q^2 = \frac{1}{M} \sum_{j=1}^M (r(qj) - q\hat{r}_1)^2. \quad (7)$$

# Asymptotic Distribution of $q$ -Daily Variance

- ▲ The asymptotic limit of the expected value of  $\widehat{\sigma}_q^2$  is

$$\mathbb{E} \left( \frac{\widehat{\sigma}_q^2}{q} \right) = \frac{1}{Mq} \sum_{j=1}^M \mathbb{E} \left( (r(qj) - q\widehat{r}_1)^2 \right) \rightarrow \sigma^2;$$

- ▲ The asymptotic limit of the variance of  $\widehat{\sigma}_q^2$  is

$$\begin{aligned} \mathbb{V} \left( \frac{\widehat{\sigma}_q^2}{q} \right) &= \frac{1}{M^2 q^2} \mathbb{V} \left( \sum_{j=1}^M (r(qj) - q\widehat{r}_1)^2 \right) \\ &= \frac{1}{Mq^2} \mathbb{V} \left( \sum_{j=1}^q u_j^2 \right) = \frac{1}{(Mq)q} \mathbb{V} \left( \sum_{j=1}^q u_j^2 \right) \\ &\rightarrow \frac{1}{Tq} \mathbb{V} (q\sigma^2 Z^2) = \frac{q}{T} \sigma^4 \mathbb{V} (Z^2) = \frac{2q\sigma^4}{T}. \end{aligned}$$

## Null Hypothesis of Variance Ratio

- ▲ By the central limit theorem, as  $T$  becomes very large,

$$\frac{\hat{\sigma}_q^2}{q} \sim N\left(\sigma^2, \frac{2q\sigma^4}{T}\right)$$

- ▲ Equivalently

$$\sqrt{T} \left( \frac{\hat{\sigma}_q^2}{q} - \sigma^2 \right) \sim N(0, 2q\sigma^4),$$

or

$$\sqrt{T} \left( \frac{\hat{\sigma}_q^2}{q\sigma^2} - 1 \right) \sim N(0, 2q).$$

- ▲ Note that the variance for the statistic  $\sqrt{T} \left( \frac{\hat{\sigma}_q^2}{q\sigma^2} - 1 \right)$ , i.e.,  $2q$ , is known!

# Test Statistics

- ▼ To perform the test, we define the test statistics

$$J_d(q) := \frac{\widehat{\sigma}_q^2}{q} - \widehat{\sigma}_1^2;$$

$$J_r(q) := \frac{\widehat{\sigma}_q^2}{q\widehat{\sigma}_1^2} - 1 = \widehat{\text{VR}}(q) - 1. \quad (8)$$

- ▼ Note that  $J_r(q) = \frac{J_d(q)}{\widehat{\sigma}_1^2}$ .

# Asymptotic Distributions

## Theorem

The asymptotic distributions of  $\sqrt{T}J_d(q)$  and  $\sqrt{T}J_r(q)$  are normal with mean 0 and variances of, respectively,  $2(q-1)\sigma^4$  and  $2(q-1)$ :


$$\sqrt{T}J_d(q) \sim N(0, 2(q-1)\sigma^4);$$

$$\sqrt{T}J_r(q) \sim N(0, 2(q-1)).$$

◆ In light of this theorem, for  $q > 1$ , the  $z$  score is computed as

$$Z_q = \sqrt{T} \frac{J_r(q)}{\sqrt{2(q-1)}} \sim N(0, 1). \quad (9)$$

# A Variance Ratio Test Algorithm



## Case Study: Variance Tests on GE

$q$	1	2	3	4	5	6	7	8	9	10
Obs	22,776	11,388	7,592	5,694	4,555	3,796	3,253	2,847	2,530	2,277
$\widehat{\text{VR}}(q)$	1	1.002	0.946	0.939	0.916	0.926	0.968	0.933	0.871	0.920
$Z_q$	—	0.20	-4.08	-3.74	-4.46	-3.53	-1.40	-2.69	-4.85	-2.86

**Table:** Results of variance ratio tests based on GE's daily log returns.


- Looking at  $Z_2$ , what can you infer?
- Looking at  $Z_5$ , what can you infer?

## Takeaways

- ✎ The Law of Iterated Expectations provides the connection between the unpredictability of return and the fundamental value of the asset.
- ✎ The variance of random walk's sample paths increases with time.
- ✎ Variance ratio test is an algorithm to test whether the asset price is a random walk.
- ✎ Empirical evidence suggests that asset prices are generally not random walks. They are not 100% random.
- ✎ Statistical arbitrage is difficult but possible because prices are not strictly random walks.
- ✎ If you invest in obtaining and analyzing financial information, it is possible to “beat the market” (though difficult).
- ✎ Smart beta based on  $VR(q)$ , skewness, and kurtosis estimates?

## Additional Reading

### 1 **Quants and the quirks: Is efficient-market theory becoming more efficient?**

Listen by clicking 

### 2 **A Non-Random Walk Down Wall Street** (Page 3 to Page 83)